

BEFORE /AFTER
BARIATRIC
SURGERY



Treatment of
12 x 45' sessions



Reduction of
visceral fat



Diminution of
health risks

Bariatric surgeons • Obesity specialists



BEFORE A BARIATRIC SURGERY:
Recommended protocol
12 sessions of 45 minutes (Daily)



Illustration of excess of abdominal fat



AFTER A BARIATRIC SURGERY:
Recommended protocol
6 sessions of 45 minutes (2 times per week)
2 weeks after the surgery



Illustration of fatty liver

DECREASE VISCERAL FAT AND LIVER SIZE



DOUBLE-BLIND RANDOMIZED CLINICAL STUDY (2010)

Dr Ghislaine Beilin, Dr Rodi Courie
Nutrition Unit - Endocrinologist - Pitié Salpêtrière Hospital

Evaluation of the average centimeter waist size reduction in **28 patients** with abdominal obesity after 12 sessions of ReduStim: **-6.1 cm.**

The other 4 metabolic syndromes criteria remain unchanged.

Fall of transaminases -23% ALAT & -13% ASAT in proportion to the drop weight and abdominal fat.

STUDY CONCLUSION:

Fatty acids released from the excess fat are not stored in the liver but completely eliminated in CO2 and H2O.